

Published in final edited form as:

Organometallics. 2010 September 13; 29(17): 3765–3768. doi:10.1021/om100262x.

Synthesis and Structure of Fused *N*-Heterocyclic Carbenes and Their Rhodium Complexes

Jean Li, Ian C. Stewart, and Robert H. Grubbs

The Arnold and Mabel Beckman Laboratories of Chemical Synthesis, Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125

New procedures for the synthesis of *N*-heterocyclic carbenes with multiple fused rings have been developed utilizing a key ring-closing metathesis step. Rhodium complexes were obtained via the pentafluorophenyl carbene adducts. Solid-state structural behavior of the new carbene ligands were analyzed via X-ray crystallography.

N-heterocyclic carbenes (NHCs), first isolated by Arduengo et al. in 1991, have become a well-studied and well-utilized class of ligands in the field of transition metal-catalyzed reactions.¹ Their proficiency in catalysis, variable electronic properties, and ease of translation into complex architectural structures have allowed for the development of unique reactivity and targeted selectivity.^{2–6} In the past 20 years, NHCs have been applied as ligands in reactions such as palladium catalyzed cross-coupling, olefin metathesis, asymmetric hydrosilylation and conjugate addition.^{7–12} Free NHCs have also shown great efficacy as organocatalysts in a variety of reactions.^{13,14} Exploration of novel structural motifs has enabled the development of novel applications for NHCs. Serving as a positive feedback loop, as the synthetic and catalytic applications have grown, the array of structural motifs has grown in parallel (Figure 1), illustrating the dynamic interchange between structure and function.^{1,15,16}

In particular, fused NHC structures are of interest, as it has been shown computationally that the rotational lability of the carbene can greatly influence the behavior and dynamics of the NHC-bound metal complex.^{17,18} In this study, the syntheses of fused carbenes **6** and **7** were designed to allow for control of stereochemistry at the backbone of the NHC as well as facile modification of the *N*-bound arene fragment. A three-carbon chain from the backbone of the NHC to the aryl was chosen to form the *cis*-fused as well as the *trans*-fused structures (Figure 2). Independent synthesis of each carbene precursor allowed for separate *cis* and *trans* routes, bypassing a need to separate the meso from the racemic forms. Synthesis of the racemate could be easily modified to render the enantioenriched mixtures of the *trans* species via resolution with inexpensive L-(+)-tartaric acid.¹⁹

Hermann and Blechert have synthesized similar NHC structures.^{16,20,21} Carbene **4** was appended to ruthenium and shown to have limited metathesis activity.¹⁶ Unfortunately, the synthesis of **4** yielded a 3:1 mixture of the meso and racemic forms, and only the meso form was isolated for study.¹⁶ The crystal structure of carbene **4**, which contains a C₂ linkage, exhibited planarity of the *N*-bound arene with the heterocycle, thereby rendering the ruthenium species noncanonical with standard ruthenium-based catalysts.

rhg@caltech.edu.

Supporting Information Available. Crystallographic data for complexes **17** and **18**, experimental procedures and NMR data for all new compounds are available.

In designing both syntheses, we chose to utilize a ring closing metathesis as the key step in fusing the seven-membered ring fragment. Previous syntheses of fused NHC frameworks began from quinoline-type derivatives and were thus limited in both ring-size and stereocontrol.¹⁶ With intimate knowledge of the olefin metathesis reaction at hand, we imagined separate syntheses hinging upon retention of the initially introduced backbone stereochemistry during ring closing metathesis (RCM).

The cis-cyclohexenyldiamine was synthesized by a known procedure.²² Palladium catalyzed cross-coupling of the free diamine with *o*-bromostyrene yielded secondary diamine **8**. Attempts at ring-closing metathesis of **8** resulted in decomposition of both diamine and the ruthenium benzylidene species. In order to cloak the diamino functionality, **8** was condensed with pentafluorobenzaldehyde to form **9**, following a recent report.²³ Ring-closing with ruthenium catalyst provided cis-fused adduct **10** and hydrogenation gave the desired carbene precursor **11**.

Entry into the racemic trans-fused carbene began with aza-Claisen rearrangement of commercially available *N*-allylaniline. Condensation with glyoxal yielded the bisimine, which was further functionalized via Grignard addition and in situ urea formation to give intermediate **12**. Although the pentafluorophenyl adduct could be introduced immediately succeeding addition of Grignard, the urea provided higher yields in the metathesis reaction, yielding ring-closed **13**.

Hydrogenation with Pd/C at 1 atm gave the urea **14**, which was readily cleaved with lithium aluminum hydride to afford trans-fused diamine **15**. The free diamine was then condensed with pentafluorobenzaldehyde to provide the carbene precursor **16**.

Recently, we reported the facile decomposition of pentafluorophenyl adducts as NHC-precursors for ruthenium complexes.²³ We chose to isolate the masked carbene as pentafluorophenyl adducts rather than the traditional imidazolium salts in order to allow ligation onto metal fragments in the absence of a strong base. Thus, reaction of **11** or **16** with rhodium cyclooctadiene chloride dimer gave the corresponding rhodium complexes shown (Scheme 3). Crystals suitable for X-ray diffraction studies were obtained as bright yellow blocks by vapor diffusion of pentane into a benzene solution of either **17** or **18**.

Figure 3 shows the ORTEP diagram of the cis-fused rhodium complex **17** obtained from X-Ray diffraction. We were surprised to note that, although there exists a cis-orientation at the backbone of the NHC, a distortion in the carbene allows the two linkers to obtain a relative trans configuration of the N-bound arenes. Note the dihedral torsion angle between C(12)-C(2)-C(3)-C(13) at 37.57° confirms a cis-orientation (Table 1). Neither arene is orthogonal to the plane of the NHC, containing dihedral angles at C(5)-C(4)-N(1)-C(1) of 67.86° and at C(20)-C(21)-N(2)-C(1) of 31.84°.

X-ray crystal analysis of trans-fused **18** confirmed the expected anti-anti relationship. Analogous to previously reported nonfused ligated NHCs, Figure 4 shows the NHC to be planar, without the distortion noted in Figure 3.

Table 2 lists selected bond lengths and angles for **18**. Here the dihedral torsion angle at the backbone of the carbene is greater than 90°, with the angle at C(12)-C(2)-C(3)-C(13) of 136.83°. Again, neither arene ring is orthogonal or planar to the NHC, with dihedral angles at C(5)-C(4)-N(1)-C(1) of 50.98° and at C(20)-C(21)-N(2)-C(1) of 62.57°.

A simplified side view of both cis and trans fused Rh-species with hydrogens omitted for clarity is shown in Figure 5. It is clear that the cis linkage exists for complex **17**, however,

the arenes are locked in a relative trans conformation, similar to that of complex **18**, which shows a higher degree of overall symmetry.

In summary, we have developed the synthesis of stereochemically defined fused *N*-heterocyclic carbene structures which can be obtained in gram quantities either as the meso form or the racemate. X-ray crystallography data confirmed the covalent linkages.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors wish to thank the NIH (F32GM078734 and K99GM084302 to I. C. S. and 5RO1GM31332 to R. H. G.) for generous financial support. J. L. thanks the National Science Foundation for a Graduate Research Fellowship. This research was made with Government support under and awarded by DoD, Air Force Office of Scientific Research, National Defense Science and Engineering Graduate (NDSEG) Fellowship, 32 CFR 168a. Dr. Michael W. Day and Lawrence M. Henling are thanked for crystallographic work, and Dr. David Vandervelde is thanked for NMR spectroscopy work.

References

1. Arduengo AJ, Harlow RL, Kline M. *J Am Chem Soc* 1991;113:361.
2. Dixon DA, Arduengo AJ. *J Phys Chem* 1991;95:4180.
3. Diez-Gonzalez S, Marion N, Nolan SP. *Chem Rev* 2009;109:3612. [PubMed: 19588961]
4. Corberan R, Mas-Marza E, Peris E. *Eur J Inorg Chem* 2009:1700.
5. Lee HM, Lee CC, Cheng PY. *Curr Org Chem* 2007;11:1491.
6. Herrmann WA, Schutz J, Frey GD, Herdtweck E. *Organometallics* 2006;25:2437.
7. Schwab P, France MB, Ziller JW, Grubbs RH. *Angew Chem Int Ed* 1995;34:2039.
8. Trnka TM, Grubbs RH. *Acc Chem Res* 2001;34:18. [PubMed: 11170353]
9. Herrmann WA. *Angew Chem Int Ed* 2002;41:1290.
10. Crudden CM, Allen DP. *Coord Chem Rev* 2004;248:2247.
11. Snead DR, Seo H, Hong S. *Curr Org Chem* 2008;12:1370.
12. de Fremont P, Marion N, Nolan SP. *Coord Chem Rev* 2009;253:862.
13. Enders D, Balensiefer T. *Acc Chem Res* 2004;37:534. [PubMed: 15311952]
14. Brown MK, May TL, Baxter CA, Hoveyda AH. *Angew Chem Int Ed* 2007;46:1097.
15. Berlin JM, Campbell K, Ritter T, Funk TW, Chlenov A, Grubbs RH. *Organic Letters* 2007;9:1339. [PubMed: 17343392]
16. Vehlouw K, Gessler S, Blechert S. *Angew Chem Int Ed* 2007;46:8082.
17. Clavier H, Correa A, Cavallo L, Escudero-Adan EC, Benet-Buchholz J, Slawin AMZ, Noaln SP. *Eur J Inorg Chem* 2009:1767.
18. Ragone F, Poater A, Cavallo L. *J Am Chem Soc* 2010;132:4249. [PubMed: 20205432]
19. Schanz HJ, Linseis MA, Gilheany DG. *Tetrahedron-Asymmetry* 2003;14:2763.
20. Herrmann WA, Baskakov D, Herdtweck E, Hoffmann SD, Bunlaksananusorn T, Rampf F, Rodefeld L. *Organometallics* 2006;25:2449.
21. Baskakov D, Herrmann WA, Herdtweck E, Hoffmann SD. *Organometallics* 2007;26:626.
22. Witiak DT, Rotella DP, Filippi JA, Gallucci J. *Journal of Medicinal Chemistry* 1987;30:1327. [PubMed: 3612684]
23. Blum AP, Ritter T, Grubbs RH. *Organometallics* 2007;26:2122.




Figure 1.
NHCs utilized in metal-mediated catalysis.




Figure 2.
Design of Rotationally Locked NHCs





Figure 3.
ORTEP diagram of **17** with 50% probability ellipsoids.



NIH-PA Author Manuscript






Figure 5.
Side-view of structures **17** and **18**.



Scheme 1.
Synthesis of cis carbene precursor **11**.



Scheme 2.
Synthesis of trans carbene precursor **16**.




Table 1Selected Bond Lengths [\AA] and Angles [$^\circ$] for **17**.

Rh(1)-C(1)	1.992(2)	C(1)-N(1)-C(2)	111.93(17)
Rh(1)-C(29)	2.105(2)	C(1)-N(2)-C(3)	110.99(17)
Rh(1)-C(22)	2.124(2)	N(1)-C(1)-N(2)	106.57(19)
Rh(1)-C(25)	2.207(2)	N(1)-C(1)-Rh(1)	124.58(14)
Rh(1)-C(26)	2.247(2)	N(2)-C(1)-Rh(1)	128.82(15)
Rh(1)-Cl(1)	2.4075(5)	N(1)-C(2)-C(3)	98.93(17)
		N(2)-C(3)-C(2)	100.04(18)
		C(12)-C(2)-C(3)-C(13)	37.57
		C(5)-C(4)-N(1)-C(1)	67.86
		C(20)-C(21)-N(2)-C(1)	31.84

Table 2Selected Bond Lengths [\AA] and Angles [$^{\circ}$] for **18**.

Rh(1)-C(1)	2.0032(7)	C(1)-N(1)-C(2)	113.12(5)
Rh(1)-C(22)	2.0969(7)	C(1)-N(2)-C(3)	113.17(5)
Rh(1)-C(29)	2.1263(8)	N(1)-C(1)-N(2)	107.40(6)
Rh(1)-C(26)	2.1847(8)	N(1)-C(1)-Rh(1)	123.40(5)
Rh(1)-C(25)	2.2169(8)	N(2)-C(1)-Rh(1)	129.10(5)
Rh(1)-Cl(1)	2.3896(2)	N(1)-C(2)-C(3)	101.40(6)
		N(2)-C(3)-C(2)	101.27(5)
		C(12)-C(2)-C(3)-C(13)	136.83
		C(5)-C(4)-N(1)-C(1)	50.98
		C(20)-C(21)-N(2)-C(1)	62.57